



Impact of processing of red beet on betalain content and antioxidant activity

Kavitha Ravichandran ^{a,*}, Nay Min Min Thaw Saw ^a, Adel A.A. Mohdaly ^a, Ahmed M.M. Gabr ^a, Anja Kastell ^a, Heidi Riedel ^a, Zhenzhen Cai ^a, Dietrich Knorr ^b, Iryna Smetanska ^{a,c}

^a Department of Methods in Food Biotechnology, Institute of Food Technology and Food Chemistry, Berlin University of Technology, Königin-Luise Str 22, 14195 Berlin, Germany

^b Department of Food Biotechnology and Food Process Engineering, Institute of Food Technology and Food Chemistry, Berlin University of Technology, Königin-Luise Str 22, 14195 Berlin, Germany

^c Department of Plant Food Processing, Agricultural Faculty, University of Applied Science Weihenstephan-Triesdorf, Steingruber Str. 2, 91746 Weidenbach, Germany

ARTICLE INFO

Article history:

Received 15 December 2010

Accepted 2 July 2011

Keywords:

Red beet

Beta vulgaris

Processing

Antioxidant activity

ABSTRACT

Red beet has high concentration betalains that are used as food colorants and food additives due to their health promoting properties. Redbeet is generally processed before consumption which influences the stability of betalains in turn which affects the acceptability and health properties. The objective of the study was to investigate the influence of processing techniques as microwaving, boiling, roasting and vacuuming on the red beet. The impact of processing was evaluated on the basis of betalains content and antioxidant activity of the processed samples. With spectrophotometric results betalains content were found to add up to 20% with vacuum treatment and also with microwave treatment of 900 W and 1800 W for 30 s up to 7% and 19% respectively, however there was decrease in boiling and roasting treatments. With HPLC analyses the content of betanin seems to increase in microwave treatments with 450 W and 900 W but reduced with 1800 W. In the case of antioxidant activity there was 2 to 3 fold increase in boiling, roasting and microwave treatments as compared to the control.

© 2011 Elsevier Ltd. All rights reserved.

1. Introduction

Colors are important quality indicators that determine the consumer acceptance of foods. In recent days market for application of synthetic colorants has decreased in favor of natural colorants (Fletcher, 2006). Fruits and vegetables are good sources of natural colorants. However, the natural colors have disadvantages as higher cost and reduced stability (Herbach, Stintzing, & Carle, 2006). It is a common practice that food from plant origin is generally processed by thermal methods to extend their shelf life, to improve flavor and to enhance the palatability. Due to processing and storage there are significant changes in betalains. According to Howard (2008) in the recent times consumption of processed foods have been increasing in greater amounts along with fresh fruits and vegetables. Betalains are water-soluble nitrogen-containing pigments, found in high concentrations in red beet (*Beta vulgaris*). Betalains consist of two sub-classes: betacyanins (red-violet pigments) and betaxanthins (yellow-orange pigments) (Delgado-Vargas, Jiménez, & Paredes-López, 2000; Stintzing & Carle, 2004). They have antimicrobial and antiviral effects (Strack, Vogt, & Schliemann, 2003) and also can inhibit the cell proliferation of human tumor cells (Reddy, Ruby, Lindo, & Nair, 2005).

Betalains can be used as food additives which either avoid the food discoloration or to enrich food. The use of betalains as food colorant is approved by European Union and betalains are labeled as E-162. Betalains are particularly suited for use colouring food products (Von Elbe, Maing, & Amundson, 1974) (Cai, Sun, Schliemann, & Corke, 2001). (Roy, Gullapalli, U.R., & R., 2004) Although anthocyanins are the most wide spread and mostly used natural pigments covering the red purple color range, betalains are more stable to pH and temperature. Betalains exhibit broad pH stability which are suited for low-acid foods where coloring with anthocyanins usually not possible. (Stintzing & Carle, 2004) For the yellow orange color range carotenoids are the natural pigments but due to poor solubility in water, betaxanthin could be used in application as yellow orange food colorants in situations (Azeredo, 2009). Betalain pigment mixtures can be used as a natural additive for food, drugs and cosmetic products in the form of beet juice concentrate or beet powder (Dörnenburg & Knorr, 1996). Consumption of red beet which are rich source of antioxidants can contribute to protection from age-related diseases. According to Vinson, Hao, Su, and Zubik (1998) Žitňanová et al. (2006) red beet is one of the most potent vegetables with respect to antioxidant activity. Betacyanins are a group of compounds exhibiting antioxidant and radical-scavenging activities (Escribano, Pedreño, García-Carmona, & Muñoz, 1998; Pedreño & Escribano, 2000). They also inhibit cervical ovarian and bladder cancer cells in vitro (Zou et al., 2005). Red beet also can be used as antioxidants (Georgiev et al., 2010). Netzel et al. (2005) reported that the ingestion of a single dose of red beet juice resulted in an increase of antioxidant compounds

Abbreviations: AOA, antioxidant activity; MW, microwave; PV, vacuum.

* Corresponding author. Tel.: +49 30 314 71263; fax: +49 30 31471492.

E-mail address: kavisri9@googlemail.com (K. Ravichandran).

including betalains in urinary excretion. Betalains and other phenolic compounds presented in red beet decreases oxidative damage of lipids and improves antioxidant status in humans. Antioxidant activity in red beet is associated involvement of antioxidants in the scavenging of free radicals and consequently in the prevention of diseases like cancer, cardiovascular diseases (Delgado-Vargas et al., 2000). Antioxidant activity was also reported to enrich human low-density lipoproteins by betalains which increase resistance to oxidation (Tesoriere, Allegra, Butera, & Livrea, 2004). According to Gentile, Tesoriere, Allegra, Livrea, and Alessio (2004) betalains exhibit anti-inflammatory effects, antiradical and antioxidant activity.

According to Stintzing and Carle (2006) maximizing betalain yield during extraction and processing is prerequisite. Extracted betalains are susceptible to pH, water activity, and exposure to light, oxygen, metal ions, temperature and enzymatic activities. However temperature is the most decisive factor for betalain decomposition within the optimal pH. However, processing changes the content of betalains and consequently food color as well as the antioxidant activity. The susceptibility of betalains to the above mentioned factors restricts their use as food colorants. The aim of our study was to investigate the influence of thermal processes as microwave, boiling, roasting and vacuuming, on the betalain content and antioxidant activity of the red beet.

2. Material and methods

2.1. Plant materials

The fresh red beet plants were obtained from the local vegetable market in Berlin, Germany and stored at 4 °C. The experiments were generally performed immediately after procurement.

2.2. Sample preparation

The red beet roots were washed and chopped into small pieces using magic mix, mechanical chopper.

For microwave irradiation was done with Microwave-Panasonic NE-1846, each sample of 10 g was placed in microwavable dishes and they were treated at 450, 900 and 1800 W for 10, 20, and 30 s.

In case of boiling 50 g of sample was treated at medium direct heat at 80 °C over a pan and stirred frequently at 30 rpm. An equal quantity of water was added to samples and heated for 60, 120, 180 sec variations.

With roasting 50 g of sample was treated at medium direct heat over a pan and stirred frequently at 30 rpm for uniform heat distribution for 60, 120, 180 s variations.

For vacuum treatment Komet plus Vacu 23 was used, about 30 g sample was packed in polypropylene bags and vacuumed at medium 94% and high 99% vacuum conditions.

All the samples after the respective treatment were cooled to room temperature, homogenized and freeze dried. The dried samples were taken for further extraction studies. All the experiments were done in triplicates.

2.3. Extraction of betalains

0.1 g of freeze dried samples were dissolved in 10 ml of 50% ethanol, were agitated for 10 s and the homogenate was centrifuged at 6000 rpm for 10 min. The supernatant was collected as it is after centrifugation and the same was repeated for 2 more times to ensure maximum extraction of betalains. The supernatant was further used for determination of betalains.

2.4. Determination of betalain compounds with spectrophotometric analyses

The content of betaxanthins and betacyanins in the extracts was determined spectrophotometrically at 538 nm and 480 nm with a UV-Vis spectrometer, respectively according to the methods of

Stintzing, Schieber, and Carle (2003). The absorbance reading obtained was used to calculate the betalain concentration for each sample. The betalain content (BC) was calculated as $BC \text{ (mg/L)} = [(A \times DF \times MW \times 1000) / (e \times l)]$, where A is the absorption, DF the dilution factor and l the pathlength (1 cm) of the cuvette. For quantification of betacyanins and betaxanthins, the molecular weights (MW) and molar extinction coefficients (e) (MW = 550 g/mol; e = 60,000 L/mol cm in H₂O) and (MW = 308 g/mol; e = 48,000 L/mol cm in H₂O) were applied.

2.5. Determination of betalain compounds with HPLC analyses

HPLC analyses were performed on HPLC (Dionex Summit P680A HPLC-System), equipped with P680 HPLC pump, ASI-100 automated sample injector, The analytical column Merck Lichrocart 250 × 4 RP-18 e (5 μm) was operated at 30 °C. Solvents were 0.2% (v/v) formic acid in water (A) and acetonitrile (B). At a flow rate of 1 mL/min, simultaneous monitoring was performed at 538 nm and 476 nm for betalains.

2.6. Antioxidant activity of extracts

The antioxidant activity of the extracted betalain samples was determined by DPPH method. The DPPH assay (Lee et al., 2003) was utilized with some modifications. The stock reagent solution (1×10^{-3} M) was prepared by dissolving 22 mg of DPPH in 50 mL of methanol and stored at -20 °C until use. The working solution (6×10^{-5} M) was prepared by mixing 6 mL of stock solution with 100 mL of methanol to obtain an absorbance value of 0.8 ± 0.02 at 515 nm, as measured using a spectrophotometer. Extracts each of 0.1 ml were vortexed for 30 s with 3.9 ml of DPPH solution and left to react for 30 min, after which the absorbance at 515 nm was recorded. A control with no added extract was also analyzed. The DPPH solution with no added extract was analyzed as control.

Scavenging activity was calculated as follows:

$$\text{DPPH radical-scavenging activity (\%)} = \left[\frac{(A_{\text{control}} - A_{\text{sample}})}{A_{\text{control}}} \right] \times 100$$

where A is the absorbance at 515 nm.

2.7. Statistical analysis

All analyses were performed in triplicate and data reported as mean ± standard deviation (SD). Data were subjected to analysis of variance (ANOVA) ($P < 0.05$). Results were processed by Excel (Microsoft Office 2007) and SPSS Version 17.0 (SPSS Inc., Chicago, IL, USA).

3. Results and discussions

The stability of betalains at different treatment conditions were investigated. Initially plant material was macerated, freeze dried and betalains were extracted with 30%, 50%, 70% ethanol and with distilled water. Efficacy of extraction was compared with different ethanol concentrations to standardize the optimal concentration for the extraction of betalains. And 50% ethanol was found to be optimal. In the literature we have found the different methods, e.g. Delgado-Vargas et al. (2000) recommended 20 to 50% of ethanol for the complete extraction of betalains. Our results of 50% ethanol are correlating the previous studies done. In our experiments 30% ethanol showed the increased extractability, while 50% ethanol showed higher stability of betalains as compared to other concentrations. For processing experiments the solvent mixtures of 50% ethanol concentration were utilized.

3.1. Spectrophotometric analyses

Betalain stability is influenced by different internal factors such as, pigment content, pH, moisture content and external factors such as temperature, light, oxygen, which needs to be considered to ensure optimum pigment and color retention in foods (Herbach et al., 2006).

Processing of foods include thermal process such as microwaving, boiling, roasting or non thermal process such as vacuum treatment. The main disadvantage in thermal processing includes betalain degradation under high temperatures. Studies reported that increasing betalain degradation rates resulting from increasing temperatures. (Barrera, Reynoso, and Mejía, 1998). Our studies with microwaving have shown that there was an increase in the betaxanthins content at higher treatment conditions with microwave irradiations like 900 watts for 30 sec and 1800 watts for 30 seconds (Fig. 1). This result was in agreement with the result of Harivaindaran, Rebecca, & Chandran, 2008 that, as there is increase in temperature the yield of betacyanin content was increased.

De Ancos, Cano, Hernandez, and Monreal (1999) studied the effects of microwave heating on the natural colors in fruit purees. Generally there were only minor changes of the quantitative and qualitative composition of carotenoids due to microwave heating of carotenoids in papaya and anthocyanins in strawberry purees. A previous enzyme inactivation by a short heat treatment of 70 °C for 2 min of the extract is desirable, to avoid betalain enzymatic degradation (Delgado-Vargas et al., 2000). According to Strack et al., 2003 acidification improves betacyanin stability and avoids oxidation by polyphenoloxidases. Red beets have several endogenous enzymes such as β -glucosidases, polyphenoloxidases and peroxidases, which if not properly inactivated may result in betalain degradation and colour losses (Escribano, Gandía-Herrero, Caballero, & Pedreño, 2002). In our experiment with microwave treatment of red beet we obtained the similar results. The quantity of betaxanthins slightly increased by 7% during 900 W for 30 s and by 19% during 1800 W for 30 s (Fig. 1).

Vacuuming as non thermal treatment prevent the betalains from degradation. We have measured the increase in betacyanin content in extracts of vacuum-dried red beet tissues from 7 to 8.5 mg/l and betaxanthin from 5.2 to 5.8 mg/l when compared non-vacuum-treated samples. This can be attributed to the better extractability of the samples (Fig. 2). There was increase in betacyanins by 20% and in betaxanthins by 12% when compared to the control.

Vacuum treatment includes removal of available oxygen or under low oxygen levels results in decreased pigment degradation than under air atmosphere. Low oxygen levels favor the pigment to be partially recovered after degradation (Von Elbe, Schwartz, & Hildenbrand, 1981). Betalains are known to be sensitive to oxidation, which has an impact on their color stability (Herbach et al., 2006).




Fig. 1. Content of betacyanins and betaxanthins after microwave treatment (BC – betacyanin and BX – betaxanthin).

Temperature is the important factor influencing betalain stability. Previous studies done Garcia-Barrera, Reynoso, and Gonzalez de Mejia (1998) have reported betalain degradation with increase in temperatures. Roy et al. (2004) have shown that the extraction of betalains from red beet was optimal at 40 °C. Boiling and roasting led to decrease in the yield of the obtained betacyanins and betaxanthins. Herbach et al. (2006) explained that thermal treatments decreased betalains stability.

As the time for boiling treatment increased from 60, 120, 180 sec, the betacyanins quantity decreased correspondingly by 6%, 22%, 51% (Fig. 3) and for betaxanthins by 18%, 23% and 33%. According to Jiratanan & Liu, 2004 betacyanin content of beets decreased by 24%, 62% and 81% and betaxanthins by 13% 60% and 73% after treatment at 105, 115 and 125 °C for 30 mins which showed the effect of processing temperatures on betacyanins and betaxanthins.

Interestingly, in the case of betaxanthins their was about 7% increase after roasting for 5 min Von Elbe et al. (1981) found in the process of canning a total of 41% pigment loss. The sensitivity of betacyanin pigments to heat and losses during processing can be due to heat liability of these colorants. During heat processing, betanin may be degraded by isomerisation, decarboxylation or cleavage by heat or acids. (Herbach, Stintzing, & Carle, 2004)

3.2. HPLC analyses

HPLC analyses were performed on HPLC (Dionex Summit P680A HPLC-System), equipped with P680 HPLC pump, ASI-100 automated sample injector, The analytical column Merck Lichrocart 250 × 4 RP-18 e (5 μ m) was operated at 30 °C. Solvents were 0.2% (v/v) formic acid in water (A) and acetonitrile (B). At a flow rate of 1 mL/min, Simultaneous monitoring was performed at 538 nm and 476 nm for betalains.

While studying the chromatograms are shown in Fig. 4, by peak area comparison of untreated red beet and treated samples, the content of betanin seems to increase in microwave treatments with 450 watts and 900 watts and also it seems to slightly increase with vacuum treatment. As the treatment temperature increases with Microwave 1800 watts and boiling the betanin content decreased. With 1800 watts betanin content was slowly reduced up to 6%. The epimerization of betanin to isobetanin may be produced due to heat or acid when food containing betanin is heated since the balance of isobetanin to betanin grows up (Fennema, 2000). HPLC analysis showed that content of betanin and isobetanin were significant with microwave treatments, betanin content was found to add up to 30% with 450 watts up to 19% with 900 waats comparison of untreated (control). While considering vacuum treatment betanin content was found to add up to 12%. In accordance with previous findings of (Herbach et al., 2004) betanin content was increased up to 75% with samples and fresh juice for 1 hr heating.




Fig. 2. Content of betalains after vacuum treatment.




Fig. 3. Content of betacyanins and betaxanthins after boiling and roasting (BC- betacyanin and BX- betaxanthin).

3.3. DPPH· radical-scavenging activity

The antioxidant activity of red beet at different treatment conditions was evaluated applying the DPPH method. When considering the antioxidant activity with longer time of processing with microwave treatments, roasting, boiling there is increase in antioxidant activity. Processed fruits and vegetables are assumed to have lower antioxidant activity compared to raw or fresh once due to degradation of vitamin C content during processing. But in our results the antioxidant activity seems to increase. Antioxidant activity of betalains is higher than of ascorbic acid (Stintzing et al., 2005).

The increase in antioxidant activities of treated samples, shows that the antioxidant activity depends not only on the presence of betalains but also other polyphenols which could have been increased during the treatments. Treatment of samples has led to better extractability and increased antioxidant activity after treatments. Except for treatments with Fig. 5. 450 W for 20 s, 900 W for 10 s and 1800 W for 10 s and also during roasting for 1 min, there was increase in antioxidant activity up to 15% in full vacuum treatment pv1 Fig. 6.




Fig. 4. HPLC Chromatogram of red beet (*Beta Vulgaris*) treated with microwaves- (a), (b), (c), (d) vacuum treatment (e), and control (f). 1. Betanin, 2. Isobetainin, 3. Betanidin.




Fig. 5. Antioxidant activity (DPPH assay) of red beet after microwave treatments.

There were increase upto 2 folds after microwave treatments. In the case of boiling and roasting in Fig. 5, there was gradual increase and up to 3 folds increase of antioxidant activity. Antioxidant activity of red beet after treatments shows that there was slight reduction in initial treatments and with increased temperatures and time the antioxidant activity was increased. The results are in agreement with the results of Dewanto, Wu, Adom and Liu (2002), and Dewanto, Wu and Liu (2002) who found that due to thermal processing there was enhance nutritional value of tomatoes and corn by increase of total antioxidant activity. This is probably related to their different contents of vitamins, and phenolic compounds, which act synergistically but are differently sensitive to processing. Overall, the increase of antioxidant activity occurring at least in the considered conditions. Also According to Adefegha and Oboh (2009) cooking causes a significant increase in antioxidant activity in tropical green leafy vegetables (Fig. 7).

4. Conclusion

Afore mentioned studies indicate that the processing of red beet has influence on betalains and antioxidant activity. Even though there was decrease in betalains content with boiling and roasting treatments Antioxidant activity was increased in microwave, boiling, roasting and vacuum treatments. But after treatments with boiling, roasting showed decreased of betalains concentration and increase in antioxidant activity are mostly indirectly correlated after treatments. These studies with thermal processing involve optimization of such processes to validate quality of phytochemicals retention. The high antioxidant activity of redbeet after treatments found good base for development of a biosynthetic process to obtain additives with a potential for application in the food, cosmetic and pharmaceutical industry.




Fig. 6. Antioxidant activity (DPPH assay) of red beet after boiling and roasting.




Fig. 7. Antioxidant activity (DPPH assay) of red beet after vacuum treatments.

Acknowledgment

The author is very grateful to Irene Hemmerich and our other colleagues of Department of Methods in Food Biotechnology and Department of Food Biotechnology and Food Process Engineering, of Berlin University of Technology, Berlin, Germany.

References

- Adefegha, S. A., & Oboh, G. (2009). Cooking enhances the antioxidant properties of some tropical green leafy vegetables. *African Journal of Biotechnology*, 10(4), 632–639.
- Azeredo, H. M. C. (2009). Betalains: properties, sources, applications, and stability: A review. *International Journal of Food Science & Technology*, 44(12), 2365–2376.
- Barrera, F., C. Reynoso y E. Mejía. (1998). Estabilidad de betalainas extraídas del garambullo (*Myrtillocactus geometrizans*). *Food Science and Technology International*, 4, 115–120.
- Cai, Y., Sun, M., Schliemann, W., & Corke, H. (2001). Chemical stability and colorant properties of betaxanthin pigments from *Celosia argentea*. *Journal of Agriculture and Food Chemistry*, 49, 4429–4435.
- De Ancos, B., Cano, M. P., Hernandez, A., & Monreal, M. (1999). Effects of microwave heating on pigment composition and colour of fruit purees. *Journal of the Science of Food and Agriculture*, 79(5), 663–670.
- Delgado-Vargas, F., Jiménez, A. R., & Paredes-López, O. (2000). Natural pigments: Carotenoids, anthocyanins, and betalains — characteristics, biosynthesis, processing, and stability. *Critical Reviews in Food Science and Nutrition*, 40, 173–289.
- Dewanto, V., Wu, X. Z., Adom, K. K., & Liu, R. H. (2002). Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity. *Journal of Agricultural and Food Chemistry*, 50, 3010–3014.
- Dewanto, V., Wu, X. Z., & Liu, R. H. (2002). Processed sweet corn has higher antioxidant activity. *Journal of Agricultural and Food Chemistry*, 50, 4959–4964.
- Dörnenburg, H., & Knorr, D. (1996). Generation of colors and flavors in plant cell and tissue cultures. *Critical Review in Plant Sciences*, 15, 141–168.
- Escribano, J., Gandía-Herrero, F., Caballero, N., & Pedreño, M. A. (2002). Subcellular localization and isoenzyme pattern of peroxidase and polyphenol oxidase in beet root (*Beta vulgaris* L.). *Journal of Agricultural and Food Chemistry*, 50, 6123–6129.
- Escribano, J., Pedreño, M. A., García-Carmona, F., & Muñoz, R. (1998). Characterization of the antiradical activity of betalains from *Beta vulgaris* L. roots. *Phytochemical Analysis*, 9, 124–127.
- Fennema, O. R. (2000). *Química de los Alimentos* (2ª edición). España: ACRIBIA S.A. Zaragoza.
- Fletcher, A. (2006). Lycopene colorant achieves regulatory approval. foodnavigator.com/news.
- García-Barrera, F. A., Reynoso, C. R., & Gonzalez de Mejia, E. (1998). Stability of betalains extracted from garambullo (*Myrtillocactus geometrizans*). *Food Science and Technology International*, 4, 115–120.
- Gentile, C., Tesoriere, L., Allegra, M., Livrea, M. A., & Alessio, P. D. (2004). Antioxidant betalains from cactus pear (*Opuntia ficus-indica*) inhibit endothelial ICAM-1 expression. *Annals of the New York Academy of Sciences*, 1028, 481–486.
- Georgiev, V. G., Weber, J., Kneschke, E. M., Nedyalkov Denev, P., Bley, T., & Pavlov, A. I. (2010). Antioxidant activity and phenolic content of betalain extracts from intact plants and hairy root cultures of the red beetroot *Beta vulgaris* cv. detroit dark red. *Plant Foods for Human Nutrition*, 65, 105–111.
- Harivandaran, K. V., Rebecca, O. P. S., & Chandran, S. (2008). Study of optimal temperature, pH and stability of dragon fruit (*Hylocereus polyrhizus*) peel for use as potential natural colorant. *Pakistan Journal of Biological Sciences*, 11(18), 2259–2263.
- Herbach, K. M., Stintzing, F. C., & Carle, R. (2004). Impact of thermal treatment on color and pigment pattern of red beet (*Beta vulgaris* L.) preparations. *Journal of Food Science*, 69, C491–C498.
- Herbach, K. M., Stintzing, F. C., & Carle, R. (2006). Betalain stability and degradation — Structural and chromatic aspects. *Journal of Food Science*, 71, R41–R50.

- Howard, L. (2008). Processing techniques and their effect on fruit and vegetable phytochemicals, improving the health-promoting properties of fruit and vegetable products. In F. A. Tomás-Barberán, & M. I. Gil (Eds.), *Woodhead publishing series in food science, technology and nutrition*, No. 157, Spain: CEBAS (CSIC).
- Jiratanan, T., & Liu, R. H. (2004). Antioxidant activity of processed table beets (*Beta vulgaris* var. *conditiva*) and green beans (*Phaseolus vulgaris* L.). *Journal of Agricultural and Food Chemistry*, 52, 2659–2670.
- Lee, S. C., Kim, J. H., Jeong, S. M., Kim, D. R., Ha, J. U., & Nam, K. C. (2003). Effect of far infrared radiation on the antioxidant activity of rice hulls. *Journal of Agricultural and Food Chemistry*, 51, 4400–4403.
- Netzel, M., Stintzing, F. C., Quaas, D., Strass, G., Carle, R., Bitsch, R., et al. (2005). Renal excretion of antioxidative constituents from red beet in humans. *Food Research International*, 38, 1051–1058.
- Pedreno, M. A., & Escribano, J. (2000). Studying the oxidation and antiradical activity of betalain from beetroot. *Journal of Biological Education*, 35, 49–59.
- Reddy, K. M., Ruby, L., Lindo, A., & Nair, G. M. (2005). Relative inhibition of lipid peroxidation, cyclooxygenase enzymes and human tumor cells proliferation by natural food color. *Journal of Agricultural and Food Chemistry*, 53, 9268–9273.
- Roy, K., Gullapalli, S., Chaudhuri, U. R., & Chakraborty, R. (2004). The use of a natural colorant based on betalain in the manufacture of sweet products in India. *International Journal of Food Science and Technology*, 39, 1087–1091.
- Stintzing, F. C., & Carle, R. (2004). Functional properties of anthocyanins and betalains in plants, food and in human nutrition. *Trends in Food Science and Technology*, 15, 19–38.
- Stintzing, F. C., & Carle, R. (2006). Cactus fruits - more than colour. *Fruit Processing*, 16, 166–171.
- Stintzing, F. C., Schieber, A., & Carle, R. (2003). Evaluation of colour properties and chemical quality parameters of cactus juices. *European Food Research and Technology*, 216, 303–311.
- Stintzing, F. C., Herbach, K. M., Mosshammer, M. R., Carle, R., Yi, W., Sellappan, S., et al. (2005). Color, betalain pattern, and antioxidant properties of cactus pear (*Opuntia* spp.) clones. *Journal of Agricultural and Food Chemistry*, 53, 442–451.
- Strack, D., Vogt, T., & Schliemann, W. (2003). Recent advances in betalain research. *Phytochemistry*, 62, 247–269.
- Tesoriere, L., Allegra, M., Butera, D., & Livrea, M. A. (2004). Absorption, excretion, and distribution of dietary antioxidant betalains in LDLs Potential health effects of betalains in humans. *The American Journal of Clinical Nutrition*, 80, 941–945.
- Vinson, J. A., Hao, Y., Su, X., & Zubik, L. (1998). Phenol antioxidant quantity and quality in foods: Vegetables. *Journal of Agricultural and Food Chemistry*, 46, 3630–3634.
- Von Elbe, J. H., Maing, I., & Amundson, C. H. (1974). Colour stability of betanin. *Journal of Food Science*, 39, 334–337.
- Von Elbe, J. H., Schwartz, S. J., & Hildenbrand, B. E. (1981). Loss and regeneration of betacyanin pigments during processing of redbeet. *Journal of Food Science*, 39, 334–337.
- Žitňanová, I., Ranostajová, S., Sobotová, H., Demelová, D., Pecháň, I., & Ďuračková, Z. (2006). Antioxidative activity of selected fruits and vegetables. *Biologia*, 61, 279–284.
- Zou, D., Brewer, M., Garcia, F., Feugang, J. M., Wang, J., Zang, R., et al. (2005). Cactus pear: A natural product in cancer chemoprevention. *Nutrition Journal*, 4, 25.